

Autism Spectrum Disorder (ASD) & Special Needs

Food reactivity patterns in complex neurodevelopmental presentations



Children with autism spectrum disorder and related special-needs presentations commonly exhibit overlapping gastrointestinal, immune, behavioral, and sensory features that can complicate nutrition planning and elimination strategies.^[1,2,6,8] IgG Food MAP may help clinicians contextualize whether recurrent exposure to commonly reactive foods aligns with gut-barrier disruption, food-specific immune activation, and certain symptom patterns.^[5,7,9]



When Might Providers Consider the IgG Food Map?

Persistent GI complaints: Constipation, diarrhea, bloating, abdominal discomfort, or mixed bowel patterns in patients with neurodevelopmental concerns have been associated with gut-brain axis disruption and higher immune-food reactivity signals.^[1,2,9]

Restricted eating patterns: Limited food variety, repetitive food preferences, or difficulty broadening diet may make it harder to identify whether commonly consumed foods are contributing to symptom burden.^[1,7]

Behavioral variability with meals: Caregiver reports of post-prandial irritability, stereotyped behaviors, or fluctuating focus may warrant a more individualized look at food-related immune associations.^[8]

History suggestive of barrier dysfunction: Eczema, recurrent inflammation, or concern for intestinal permeability may increase interest in assessing foods linked in the literature with intestinal permeability marker.^[5,6,15]

Trial-and-error elimination fatigue: When broad gluten-free/casein-free or other restriction strategies have produced unclear results, individualized reactivity patterns may better support structured clinical reasoning.^[12,13,14,15]

Coexisting dysbiosis or yeast concerns: GI dysfunction with suspected microbial imbalance or Candida-related burden may justify a broader review of food and yeast-directed immune exposures.^[10,11]

What the IgG Food MAP Measures

Analyte Class	What it Reflects	ASD & Special Needs Relevance
Wheat / gluten-related proteins	Bread, pasta, crackers, cereals, baked foods, and gluten-containing snacks are routine exposure sources for gliadin and related wheat antigens. ^[3]	Gliadin- and wheat-related IgG findings have been reported more often in ASD cohorts and may help contextualize gut-barrier and immune activation patterns. ^[3,4,5]
Dairy proteins	Milk, cheese, yogurt, whey-containing foods, casein-containing products, and many processed snacks provide ongoing exposure to cow's milk proteins. ^[5,9]	Dairy-directed IgG responses are frequently discussed alongside intestinal permeability biomarkers and elimination-diet decision-making in this population. ^[5,9,17]

What the IgG Food MAP Measures (Cont)

Analyte Class	What it Reflects	ASD & Special Needs Relevance
Egg proteins	Eggs in breakfasts, baked goods, sauces, breads, and packaged foods can create frequent low-visibility exposure. ^[5]	Egg-related IgG has been associated with intestinal permeability marker patterns and may help frame delayed immune reactivity in patients with mixed GI and behavioral complaints. ^[5]
Broad mixed food panel	Daily exposure can include legumes, grains, meats, fruits, vegetables, nuts, and other staple foods that vary by diet pattern. ^[8,9]	Broad food-specific IgG positivity has been reported in pediatric ASD cohorts, supporting individualized review rather than assumption based on only one or two foods. ^[8,9]
Yeast-related antigens	Candida albicans and Saccharomyces-related exposures may intersect with dysbiosis, fermentation, and yeast-associated immune responses. ^[10,11]	Yeast-directed IgG may be relevant when ASD presentations include GI dysfunction, suspected dysbiosis, or immune activation beyond standard food proteins. ^[10,11]

Related Conditions

Condition	Clinical Encapsulation
Autism spectrum disorder	Published literature describes elevated food-specific IgG, altered gut microbiota, and barrier dysfunction in subsets of ASD patients, supporting mechanistic plausibility rather than diagnostic confirmation. ^[1,2,3,6,8,9]
GI complaints	Constipation, diarrhea, bloating, and abdominal discomfort commonly co-occur with ASD and may intersect with dysbiosis, permeability changes, and food-immune signaling. ^[1,2,6]
Stereotyped or behavioral symptoms	Higher food-specific IgG concentrations have been correlated with stereotyped behavior severity in observational pediatric ASD data, although directionality remains unresolved. ^[8]
Atopic dermatitis / eczema	Food-specific IgG4-guided elimination has shown symptom improvement in pediatric allergic presentations, offering adjacent evidence context for immune-guided nutrition strategies. ^[15]
Immune dysregulation / barrier concerns	Wheat, dairy, and egg IgG associations with intestinal permeability markers suggest a possible link between food reactivity patterns and compromised epithelial integrity. ^[5]
Yeast-associated dysbiosis concerns	Anti-Candida IgG positivity and higher colonization frequency reported in ASD literature suggest yeast-related immune burden may coexist with food-reactivity patterns in some patients. ^[10,11]

How the IgG Food MAP Supports Clinical Reasoning

Individualizes elimination planning: Shifts the conversation from broad empiric restriction toward patient-specific food reactivity patterns that may guide a more structured trial. ^[14,15]	Adds gut-immune context: May help clinicians interpret nutrition-related symptoms alongside literature on permeability, immune complexes, cytokine signaling, and microbiome disruption. ^[1,5,6,16]	Supports pattern recognition: Can help connect commonly reactive foods such as wheat, dairy, and egg with recurring GI, skin, or behavioral observations documented by caregivers. ^[3,5,8]	Frames follow-up decisions: Provides one data point that may be integrated with diet history, symptom response, stool or microbiome findings, and broader functional assessment. ^[2,6,10]
---	--	---	--

How to Order

To order the IgG Food Map test, contact your Mosaic Diagnostics representative or visit [www.mosaicdx.com]. Specimen collection kits and requisition forms are available through your account portal. For clinical support or interpretation questions, contact our clinical education team.

This document is intended for licensed healthcare professionals and is for informational and educational purposes only. The test results are not intended to diagnose, treat, cure, or prevent disease. All clinical interpretation should be done by a qualified healthcare professional in the context of the individual patient. Statements regarding associations between health symptoms and conditions do not constitute diagnostic claims.

References

1. Li H, et al. Autism spectrum disorder and eating problems: imbalance of the gut-brain axis. *Nutrients*. 2021.
2. Vuong HE, Hsiao EY. The gut-microbiota-brain axis in autism spectrum disorder. *NCBI Bookshelf/StatPearls*. 2021.
3. Dettmer K, et al. Antibodies against food antigens in patients with autistic spectrum disorders. *BioMed Research International*. 2013.
4. Sanctuary MR, et al. Children with elevated wheat IgG4 antibody titer in autism spectrum disorder. *Pediatric Allergy and Immunology*. 2019.
5. Vojdani A, et al. Associations between food-specific IgG antibodies and intestinal permeability biomarkers. *Frontiers in Nutrition*. 2022.
6. Frye RE, et al. Leaky gut plays a critical role in the pathophysiology of autism. *Frontiers in Immunology*. 2022.
7. Careaga M, et al. Increased IgG4 levels in children with autism disorder. *Brain, Behavior, and Immunity*. 2008.
8. Guo M, et al. Study on aberrant eating behaviors, food intolerance, and stereotyped behaviors in autism spectrum disorder. *Frontiers in Psychiatry*. 2020.
9. Li C, Liu Y, Fang H, Chen Y, Weng J, Zhai M, Xiao T, Ke X. Study on Aberrant Eating Behaviors, Food Intolerance, and Stereotyped Behaviors in Autism Spectrum Disorder. *Front Psychiatry*. 2020 Nov 5;11:493695. doi: 10.3389/fpsy.2020.493695. PMID: 33240114; PMCID: PMC7678488.
10. Wheeler AC, et al. Anti-Candida albicans IgG antibodies in children with autism. *Journal of Child and Adolescent Psychopharmacology*. 2018.
11. Mora-Contreras AM, et al. Might Candida albicans be a risk factor for autism? A systematic review. *International Journal of Pediatrics*. 2025.
12. Gluten-free/casein-free diet trial literature summarized in the attached review, including double-blind and placebo-controlled findings.
13. González-Domenech PJ, et al. Influence of a combined gluten-free and casein-free diet on behavior in ASD. *Journal of Autism and Developmental Disorders*. 2020.
14. Gogou M, et al. Gluten-free casein-free diet for autism spectrum disorders. *Nutrients*. 2020.
15. Calderoni S, et al. Food-specific IgG4-guided diet elimination improves allergic symptoms in children. *Frontiers in Immunology*. 2024.
16. Zhang Y, et al. Food-specific IgG-based elimination diet decreased IL-6, TNF-alpha and improved symptoms. *Frontiers in Nutrition*. 2025.

Visit MosaicDX.com for more information or contact us at (800) 288-0383

All trademarks are owned by the company and its affiliates. ©2026 Mosaic Diagnostics. All Rights Reserved. MDX_ASD_IGG_09-2026.v2